

AD \_\_\_\_\_

GRANT NUMBER DAMD17-97-1-7109

TITLE: A Modulator of FGF's in Breast Cancer

PRINCIPAL INVESTIGATOR: Violaine K. Harris

CONTRACTING ORGANIZATION: Georgetown University Medical Center  
Washington, DC 20007

REPORT DATE: August 1998

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;  
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

**DTIC QUALITY INSPECTED 1**

## REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE August 1998	3. REPORT TYPE AND DATES COVERED Annual (1 Aug 97 - 31 Jul 98)
4. TITLE AND SUBTITLE A Modulator of FGF's in Breast Cancer		5. FUNDING NUMBERS DAMD17-97-1-7109
6. AUTHOR(S) Harris, Violaine K.		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Georgetown University Medical Center Washington, DC 20007		8. PERFORMING ORGANIZATION REPORT NUMBER
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012		10. SPONSORING / MONITORING AGENCY REPORT NUMBER
11. SUPPLEMENTARY NOTES		
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited		12b. DISTRIBUTION CODE  19990301008
13. ABSTRACT (Maximum 200 words) <p>The growth of new blood vessels, or angiogenesis, is an important part of breast cancer biology. We have found that a secreted binding protein for fibroblast growth factors (FGF-BP) is expressed in many breast cancer cell lines and primary breast tumor samples. In addition, FGF-BP is expressed in squamous cell carcinoma and colon cancer and modulation of FGF-BP expression in these tumor types has a significant effect on tumor growth and angiogenesis. In order to better understand the regulation of FGF-BP and how its aberrant expression might lead to activation of angiogenic pathways, I isolated the human FGF-BP promoter and determined which functional promoter elements were necessary for its expression. In particular, I found that the FGF-BP gene is transcriptionally up-regulated by the phorbol ester TPA (12-O-tetradecanoylphorbol-13-acetate) which activates the protein kinase C (PKC) pathway. The positive regulatory elements which mediate FGF-BP induction by TPA include a juxtaposed Ets/AP-1 site as well as a C/EBP site. Furthermore, the presence of a distinct repressor element was detected which normally limits the response of the promoter to TPA. Additionally, I found that the FGF-BP is induced in the presence of EGF (epidermal growth factor), which is known to play an important role in the pathogenesis of breast cancer.</p>		
14. SUBJECT TERMS Breast Cancer		15. NUMBER OF PAGES 8
		16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified
		20. LIMITATION OF ABSTRACT Unlimited

## FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.


✓ In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

✓ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

✓ In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

✓ In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

  
PI - Signature

10-13-98  
Date

## Annual report for Grant Number DAMD17-97-1-7109

P.I. Violaine K. Harris

Title: A Modulator of FGF's in Breast Cancer

**I. Introduction****Growth factors, tumor angiogenesis and metastasis**

One of the pivotal roles of locally-acting polypeptide growth factors is the induction of new blood vessels in a healing wound as well as in growing tumors. It has been shown in numerous studies with different approaches that a solid tumor cannot grow beyond a few millimeters in size without sufficient blood supply. In addition to the nourishing function of tumor blood vessels, they provide a pathway for the tumor cells to metastasize to distant organs (1-5). A direct correlation between blood vessel density in primary tumors and their metastasis has been reported for breast cancer (6-10). Most interestingly tumor angiogenesis as reflected in microvessel density is an independent prognostic indicator in breast cancer patients when tested against known parameters (e.g. tumor size, estrogen receptor, lymph node status, c-erbB-2 expression). Since angiogenesis is such an important feature of *in vivo* tumor biology, the driving forces behind this process need to be understood.

**The role of fibroblast growth factors (FGFs) and a potential function for a novel binding protein (BP) for FGF**

The most prominent and best studied angiogenesis factors are members of the fibroblast growth factor (FGF) family of polypeptides (11,12). Several members of the FGF family, mainly bFGF and aFGF are very effective angiogenesis factors and have thus been a focus of research in tumor vascularization in the past decade. The biological activities of both FGFs can be quenched by tight binding to heparansulfate proteoglycans present in the extracellular matrix (13-16). It is only partly understood how these FGFs become solubilized and thus activated in embryonic or in tumor tissues that require angiogenesis for their growth. One established mechanism that can solubilize bFGF from this storage site is the digestion of the glycosaminoglycan portion of the cell attachment molecule by heparanases (17-20).

An alternate mode of delivering active FGF from the storage site to its receptor could be binding to a secreted carrier protein. Recently, a secreted protein has been described binds to aFGF and bFGF in a non-covalent, reversible manner (21). Furthermore, bFGF bound to this protein was prevented from degradation and retained its mitogenic activity (21). These characteristics make this **FGF-binding protein (BP)** an excellent candidate carrier molecule for FGFs. We hypothesized that this BP could be an important regulator that releases immobilized FGFs from their matrix storage site and thus activates them *in vivo* (22,23).

**The secreted binding protein for FGF (BP) as a carrier for immobilized FGFs**

Recently, it was demonstrated by our laboratory that expression of this binding protein in cell lines which express bFGF leads to a tumorigenic and angiogenic phenotype of these cells (22). We also showed that BP-transfected cells release the protein into their media together with bFGF in a non-covalently bound form. This released bFGF becomes activated biologically. In addition, *in vivo* tumor growth of a BP-positive squamous cell carcinoma (SCC) cell line and a colon cancer cell line was inhibited by a reduction of endogenous BP using BP-targeted ribozymes (24). This supports the notion of an activating step for the locally stored bFGF due to expression and secretion of BP.

BP mRNA is expressed in tumor cell lines and primary tumor tissue of squamous cell carcinomas, colon cancers, and breast cancers (22). Using skin carcinogenesis as a model for epithelial cancers, we studied the role of BP expression during tumor progression. We found that BP mRNA is upregulated in the skin during development but drops to low levels in the adult mouse skin. Interestingly, in both mouse and human skin, BP mRNA and protein levels increase more than 3-fold upon treatment with the PKC-activating agent TPA (12-O-tetradecanoylphorbol-13-acetate), and is further upregulated (4-7 fold) in DMBA/TPA induced papillomas and carcinomas (25,26). The correlation between BP expression and tumor promotion by TPA strongly suggests a role for BP in tumorigenesis. The study of BP regulation in the mouse skin, which I co-authored, was published last year (25).

Using RT-PCR to screen for FGF-BP expression, we found that FGF-BP is expressed in 4 out of 6 clinical samples of primary human breast cancers, and in 9 out of 14 breast cancer cell lines. We also detected BP

mRNA expression in the mammary gland of the human and the mouse. This pattern of expression suggests that BP may be involved in breast cancer progression, where its deregulated expression could potentially contribute to tumor growth and angiogenesis. During my thesis research I plan experiments that will test the role of BP in human breast cancer cell progression, and its mechanism of regulation by the tumor promoter TPA.

The aims of my grant application were the following:

**Aim 1:** To study the tumor growth effects of BP expression in breast cancer cells. and **Aim 2:** To study the mechanisms of regulation of BP by Protein Kinase C (PKC).

## **II. Summary of Results**

The results regarding Aim 2 will be addressed in this report. This work, for which I was first author, has previously been published in the **Journal of Biological Chemistry in July 1998 (27)**. In order to investigate the mechanism of TPA induction of FGF-BP gene expression, we examined the level of FGF-BP mRNA after treatment in the ME-180 cell line, which expresses high levels of endogenous FGF-BP. We found that TPA increased FGF-BP mRNA levels in a time- and dose-dependent manner. Furthermore, induction was mediated *via* the PKC signal transduction pathway, since treating with a specific PKC inhibitor, Calphostin C, blocked the effect of TPA. Results from actinomycin D and cycloheximide experiments as well as nuclear transcription assays revealed that TPA upregulated the steady-state levels of FGF-BP mRNA by increasing its rate of gene transcription independently of *de novo* protein synthesis.

**Isolation and Characterization of the Human FGF-BP Promoter.** In order to better understand the transcriptional regulation of the human FGF-BP gene, 1.8kb of genomic sequence upstream to the known 5'UTR sequence of human FGF-BP cDNA was isolated from a human genomic library and sequenced and the transcription start site of the human gene was determined using primer extension analysis. In addition, alignment between the human and mouse FGF-BP promoter which we cloned previously (25) revealed a region of high homology with 70% nucleotide identity within the first 200 nucleotides upstream from the transcription start. Nucleotide homology dropped significantly in more upstream sequences, suggesting that the proximal conserved 200 nucleotides of the promoter could be important for transcriptional regulation of FGF-BP in both species.

Sequence analysis of the promoter demonstrated the presence of numerous consensus transcription factor binding sites which were conserved between mouse and human FGF-BP promoters and which may have functional importance in FGF-BP regulation. Consensus binding sites included TATA box is located at about -25 upstream from the transcription start for both promoters. Also, a highly conserved consensus binding site for C/EBP $\beta$ , a member of the CCAAT/enhancer binding protein (C/EBP) family of leucine zipper transcription factors which plays a central role in the acute-phase response and in a number of cell differentiation pathways (28-30). An AP-1 consensus binding site lies juxtaposed to a sequence with homology to an Ets factor binding motif, suggesting potential functional similarity to the juxtaposed Ets/AP-1 site found genes involved in invasion and metastasis such as collagenase and urokinase plasminogen activator (uPA) (31,32). In addition, a consensus Sp1 factor binding site, an additional Ets factor binding motif, and a potential NF- $\kappa$ B binding site are located in the conserved region of the promoter and may play a role in transcriptional regulation of FGF-BP as well.

**Functional analysis of the human FGF-BP promoter.** To identify the functional promoter elements involved in FGF-BP gene regulation by TPA, progressive 5' deletion mutants as well as internal deletion mutants were constructed based on the location of consensus factor binding sites on the promoter. Deletion constructs were transiently transfected into ME-180 cells and their relative luciferase activity was assayed in the absence or presence of TPA. Basal activity of the FGF-BP promoter was mediated by a combination of the Sp1 site, as well as by the juxtaposed Ets/AP-1 site.

The regulatory elements mediating TPA induction of the promoter were all contained in the first 118 base pairs upstream of the transcription start site. Further mutational analysis revealed that full TPA induction required an interplay between several regulatory elements, including the Ets/AP-1 site and the C/EBP site.

**TPA regulation of the FGF-BP promoter involves a repressor element juxtaposed to the AP-1 site.** Between the AP-1 site and the C/EBP $\beta$  site lies a region of low homology between the human and mouse BP promoter sequences. Because this region was not suspected to have any effect on TPA induction, an internal deletion removing this region (-57 to -47) was tested as a control. Surprisingly, in the  $\Delta 57/47$  construct, TPA induction of the FGF-BP promoter increased from approximately 5-fold to 11-fold, suggesting the presence of a possible repressor which may interact with this site. The lack of sequence conservation between the human

and mouse in this region may reflect a difference in the regulation of FGF-BP between the two species. The -57 to -47 deletion disrupts an AACGTG (-60 to -55) which is juxtaposed to the 3' end of the AP-1 site and which shows some similarity to the CACGTG E-box element recognized by a number of helix-loop-helix factors (33). To test this imperfect E-box for repressor activity, a C to T point mutation at position -58 was introduced into the -118/+62 BP promoter construct. The m-58 construct showed a dramatic increase in TPA induction up to 16-fold above background. This data shows that the point mutation at position -58, as well as the internal deletion from -57 to -47, disrupts repression of the FGF-BP promoter which normally limits the response to TPA.

**Transcription factor binding to FGF-BP promoter elements.** In order to ascertain that TPA-induction of FGF-BP was due to direct activation by transcription factors, we performed gel retardation analysis to show transcription factor binding to FGF-BP promoter elements. This analysis showed specific and TPA-inducible protein binding to the Ets, AP-1, and C/EBP sites. Furthermore, distinct, specific and TPA-inducible binding to the imperfect E box repressor element was also apparent. **Overall, this data indicated that TPA effects on FGF-BP gene transcription are tightly controlled by a complex interplay of positive elements and a novel negative regulatory element (27).**

**EGF induction of the FGF-BP gene.** During the course of these experiments, we found that both fetal bovine serum (FBS) and the epidermal growth factor (EGF) can regulate FGF-BP expression. Using Northern blot analysis to detect endogenous levels of FGF-BP mRNA, we found that EGF and FBS treatment causes a rapid induction of FGF-BP gene expression (unpublished results). EGF induction of FGF-BP was not dependent on new protein synthesis, suggesting that this is a direct effect on the FGF-BP gene. Furthermore, EGF and FBS induction were dependent on the PKC pathway, since treatment with a PKC inhibitor blocked their effects. Preliminary analysis of the FGF-BP promoter has shown that some of the similar transcriptional elements (i.e. Ets/AP-1 and C/EBP) are required for full induction of the promoter by EGF. Regulation of FGF-BP by EGF is highly significant to breast cancer. The EGF family of growth factors plays an important role in the development of the mammary gland and in the pathogenesis of breast cancer.

### III. Publications and Abstracts

Harris, V.K., Liaudet-Coopman, E.D.E., Boyle, B.J., Wellstein, A., Riegel, A.T. Phorbol ester-induced transcription of a FGF-binding protein is modulated by a complex interplay of positive and negative regulatory promoter elements. *J Biol Chem* 273: 19130-19139. 1998 July.

Harris, V.K., Liaudet-Coopman, E., Wellstein, A., Riegel, A.T. A fibroblast growth factor binding protein (FGF-BP) is transcriptionally regulated by phorbol esters and retinoic acid. *Proceedings of the American Association for Cancer Research*, 89th Annual Meeting, New Orleans, LA, March 28-April 1, 1998

#### **IV. Bibliography**

1. Folkman, J. and Klagsbrun, M. (1987) Angiogenic factors. *Science* **235**, 442-447
2. Cross, M. and Dexter, T. M. (1991) Growth factors in development, transformation, and tumorigenesis. *Cell* **64**, 271-280
3. Liotta, L. A.; Steeg, P. S. and Stetler-Stevenson, W. G. (1991) Cancer metastasis and angiogenesis: An imbalance of positive and negative regulation. *Cell* **64**, 327-336
4. Folkman, J. (1986) How is blood vessel growth regulated in normal and neoplastic tissue? - G.H.A. Clowes memorial award lecture. *Cancer Res.* **46**, 467-473
5. Fidler, I. F. and Ellis, L. M. (1994) The implications of angiogenesis for the biology and therapy of cancer metastasis. *Cell* **79**, 185-188
6. Horak, E. R.; Leek, R.; Klenk, N.; Lejeunde, S.; Smith, K.; Stuart, N.; Greenall, M.; Stepniewska, K. and Harris, A. L. (1992) Angiogenesis, assessed by platelet/endothelial cell adhesion molecule antibodies, as indicator of node metastases and survival in breast cancer. *Lancet* **340**, 1120-1124
7. Weidner, N.; Semple, J. P.; Welch, W. R. and Folkman, J. (1991) Tumor angiogenesis and metastasis--correlation in invasive breast carcinoma. *N. Engl. J Med.* **324**, 1-8
8. Bosari, S.; Lee, A. K.; DeLellis, R. A.; Wiley, B. D.; Heatley, G. J. and Silverman, M. L. (1992) Microvessel quantitation and prognosis in invasive breast carcinoma. *Hum. Pathol.* **23**, 755-761
9. Weidner, N.; Folkman, J.; Pozza, F.; Bevilacqua, P.; Allred, E. N.; Moore, D. H.; Meli, S. and Gasparini, G. (1992) Tumor angiogenesis: a new significant and independent prognostic indicator in early-stage breast carcinoma [see comments]. *J Natl. Cancer Inst.* **84**, 1875-1887
10. Toi, M.; Kashitani, J. and Tominaga, T. (1993) Tumor angiogenesis is an independent prognostic indicator in primary breast carcinoma. *Int. J Cancer* **55**, 371-374
11. Baird, A. and Klagsbrun, M. (1991) The fibroblast growth factor family. *Cancer Cells* **3**, 239-243
12. Gospodarowicz, D.; Ferrara, N.; Schweigerer, L. and Neufeld, G. (1987) Structural characterization and biological functions of fibroblast growth factor. *Endocr. Rev.* **8**, 95-114
13. Vlodavsky, I.; Folkman, J.; Sullivan, R.; Fridman, R.; Ishai-Michaeli, R.; Sasse, J. and Klagsbrun, M. (1987) Endothelial cell-derived basic fibroblast growth factor: synthesis and deposition into subendothelial extracellular matrix. *Proc Natl. Acad Sci. USA* **84**, 2292-2296
14. Rogelj, S.; Klagsbrun, M.; Atzmon, R.; Kurokawa, M.; Haimovitz, A.; Fuks, Z. and Vlodavsky, I. (1989) Basic fibroblast growth factor is an extracellular matrix component required for supporting the proliferation of vascular endothelial cells and the differentiation of PC12 cells. *J. Cell Biol.* **109**, 823-831
15. Saksela, O.; Moscatelli, D.; Sommer, A. and Rifkin, D. B. (1988) Endothelial cell-derived heparan sulfate binds basic fibroblast growth factor and protects it from proteolytic degradation. *J Cell Biol* **107**, 743-751
16. Kiefer, M. C.; Stephans, J. C.; Crawford, K.; Okino, K. and Barr, P. J. (1990) Ligand-affinity cloning and structure of a cell surface heparan sulfate proteoglycan that binds basic fibroblast growth factor. *Proc. Natl. Acad. Sci. U. S. A.* **87**, 6985-6989
17. Vlodavsky, I.; Eldor, A.; Bar-Ner, M.; Fridman, R.; Cohen, I. R. and Klagsbrun, M. (1988) Heparan sulfate degradation in tumor cell invasion and angiogenesis. *Adv. Exp. Med. Biol.* **233**, 201-210

18. Bashkin, P.; Doctrow, S.; Klagsbrun, M.; Svahn, C. M.; Folkman, J. and Vlodavsky, I. (1989) Basic fibroblast growth factor binds to subendothelial extracellular matrix and is released by heparitinase and heparin-like molecules. *Biochemistry*. **28**, 1737-1743
19. Moscatelli, D. (1992) Basic fibroblast growth factor (bFGF) dissociates rapidly from heparan sulfates but slowly from receptors. Implications for mechanisms of bFGF release from pericellular matrix. *J. Biol. Chem.* **267**, 25803-25809
20. Vlodavsky, I.; Bashkin, P.; Ishai-Michaeli, R.; Chajek-Shaul, T.; Bar-Shavit, R.; Haimovitz-Friedman, A.; Klagsbrun, M. and Fuks, Z. (1991) Sequestration and release of basic fibroblast growth factor. *Ann. N. Y. Acad. Sci.* **638**, 207-220
21. Wu, D.; Kan, M.; Sato, G. H.; Okamoto, T. and Sato, J. D. (1991) Characterization and molecular cloning of a putative binding protein for heparin-binding growth factors. *J. Biol. Chem.* **266**, 16778-16785
22. Czubayko, F.; Smith, R. V.; Chung, H. C. and Wellstein, A. (1994) Tumor growth and angiogenesis induced by a secreted binding protein for fibroblast growth factors. *J Biol. Chem.* **269**, 28243-28248
23. Rak, J. and Kerbel, R. S. (1997) bFGF and tumor angiogenesis-Back in the limelight? *Nature Med.* **3**, 1083-1084
24. Czubayko, F.; Liaudet-Coopman, E. D. E.; Aigner, A.; Tuveson, A. T.; Berchem, G. and Wellstein, A. (1997) A secreted FGF-binding protein can serve as the angiogenic switch for human cancer. *Nature Med.* **3**, 1137-1140
25. Kurtz, A.; Wang, H. L.; Darwiche, N.; Harris, V. and Wellstein, A. (1997) Expression of a binding protein for FGF is associated with epithelial development and skin carcinogenesis. *Oncogene*. **14**, 2671-2681
26. Aigner, A.; Kurtz, A.; Cabal, R. H.; Butler, R. E.; Hood, D. R.; Sessions, R. B.; Schulte, A.; Czubayko, F. and Wellstein, A. (1998) An FGF-binding protein during wound healing and carcinogenesis of human skin. *submitted*
27. Harris, V. K.; Liaudet-Coopman, E. D. E.; Boyle, B. J.; Wellstein, A., and Riegel, A. T. (1998) Phorbol ester-induced transcription of a FGF-binding protein is modulated by a complex interplay of positive and negative regulatory promoter elements. *J. Biol. Chem.* **273**, 19130-19139
28. Wedel, A. and Ziegler-Heitbrock, H. W. (1995) The C/EBP family of transcription factors. *Immunobiol.* **193**, 171-185
29. Akira, S. and Kishimoto, T. (1992) Il-6 and NF-IL6 in acute-phase response and viral infection. *Immunol. Rev.* **127**, 25-50
30. Cao, Z.; Umek, R. M. and McKnight, S. L. (1991) Regulated expression of three C/EBP isoforms during adipose conversion of 3T3-L1 cells. *Genes Dev.* **5**, 1538-1552
31. Gutman, A. and Wasylyk, B. (1990) The collagenase gene promoter contains a TPA and oncogene-responsive unit encompassing the PEA3 and AP-1 binding sites. *EMBO J.* **9**, 2241-2246
32. Rorth, P.; Nerlov, C.; Blasi, F. and Johnsen, M. (1990) Transcription factor PEA3 participates in the induction of urokinase plasminogen activator transcription in murine keratinocytes stimulated with epidermal growth factor and phorbol-ester. *Nucleic Acids Res.* **18**, 5009-5017
33. Kadesch, T. (1993) Consequences of heteromeric interactions among helix-loop-helix proteins. *Cell Growth Diff.* **4**, 49-55